

2016 Missouri Student Survey Dade County





2016	2016	2014
County Data	MO Data*	County Data

	•		<i>3</i>
Approximate Sample Size	201		198
Demographics (% reflects "yes" answer)			
Grade level. 6th Grade	16.5%	13.9%	12.6%
Grade level. 7th Grade	11.3%	14.4%	18.9%
Grade level. 8th Grade	15.7%	14.3%	21.1%
Grade level. 9th Grade	17.4%	15.2%	10.5%
Grade level. 10th Grade	17.4%	14.7%	16.8%
Grade level. 11th Grade	9.6%	13.9%	12.6%
Grade level. 12th Grade	12.2%	13.6%	7.4%
Male	51.3%	47.7%	51.6%
Hispanic or Latino	1.8%	2.1%	1.1%
Race, alone or in combination. Black or African Am.	0.0%	16.0%	0.0%
Race, alone or in combination. White	94.8%	78.8%	97.9%
Race, alone or in combination. Other	0.9%	1.2%	1.1%

Note: Small sample sizes at the county level can make comparisons between years difficult. Take careful note of the demographic information listed above. Drastic changes in the sample, especially changes in the percent per grade, can influence the rest of the data in this report. This means changes between years may be a result of differing samples rather than an actual population change.

^{*}Missouri data reflects a random sample, weighted to proportionally reflect the state student population.

Substances: Past 30 days (% reflects "1+" answer)			
Used alcohol	21.7%	14.2%	13.8%
Used electronic cigarettes	12.2%	10.6%	5.4%
Used Rx not prescribed for you by a doctor	8.6%	10.0%	6.4%
Used tobacco (cigarettes or chew)	17.2%	9.9%	10.8%
Used marijuana	6.9%	7.0%	5.4%
Used hookahs or water pipes	5.2%	3.4%	1.1%
Used 'OTC' drugs for non-medical reasons	1.7%	2.2%	0.0%
Used inhalants	1.7%	1.1%	1.1%
Used synthetic drugs	0.0%	0.3%	1.1%

	2016 County Data	2016 MO Data *	2014 County Data
Substances: Ever in your life (% reflects "1+" answer)			
Used hallucinogens such as LSD (acid), PCP (angel dust), or magic mushrooms	1.7%	1.1%	1.1%
Used cocaine or 'crack'	0.9%	0.9%	1.1%
Used 'club drugs' such as ecstasy	0.9%	0.8%	1.1%
Used methamphetamine (meth, crank, crystal, ice)	0.9%	0.3%	1.1%
Used heroin or 'smack'	0.0%	0.2%	0.0%
Behavior: Last 30 Days (% reflects "1+" answer)			
Missed whole day(s) of school because you skipped or cut	34.8%	29.0%	30.1%
Did not go to school because you felt you would be unsafe at school or on your way to or from school	7.8%	5.9%	5.3%
Carried a weapon such as a gun, knife or club on school property	9.6%	4.0%	7.4%
Rode with someone who was drinking alcohol	12.9%	14.3%	13.7%
Behavior: Last 3 Months (% reflects "1+" answer)			
Spread mean rumors or lies about other kids at school	28.4%	22.9%	22.3%
Posted something online or sent a text that might embarrass or hurt another student	16.7%	16.5%	14.7%
Made fun of other people	64.9%	54.1%	52.1%
Hit, shoved, or pushed another student and was not fooling around	22.6%	15.5%	18.9%
Behavior: Last 12 Months (% reflects "1+" answer)			
Been bullied on school property	38.3%	28.8%	34.4%
Were in a physical fight	28.4%	17.3%	19.1%
Been threatened or injured with a weapon such as a gun, knife or club on school property	12.2%	7.4%	7.5%
Considered suicide	13.2%	13.9%	10.5%
Planned suicide	10.4%	9.9%	8.5%
Attempted suicide	9.6%	6.3%	4.3%
Depression: Past Month (% reflects often + always)			
Was very sad	23.3%	22.7%	20.4%
Was grouchy or irritable, or in a bad mood	31.3%	33.6%	24.5%
Felt hopeless about the future	14.8%	13.5%	12.8%
Felt like not eating or eating more than usual	17.4%	21.7%	16.0%
Felt like sleeping a lot more or a lot less than usual	20.7%	24.8%	13.2%
Had difficulty concentrating on school work	31.0%	29.6%	25.5%

	2016 County Data	2016 MO Data *	2014 County Data
Perception of wrongness (% reflects wrong + very wrong))		
'Wrong' or 'very wrong' to use Rx drugs that have not been prescribed to you	95.7%	94.0%	91.3%
'Wrong' or 'very wrong' to use Cold/Cough Medicines or OTC meds to get high	95.7%	92.3%	91.4%
'Wrong' or 'very wrong' to have 5+ drinks, once or twice a week	81.7%	88.7%	87.1%
'Wrong' or 'very wrong' for you to smoke tobacco	83.5%	88.4%	89.2%
'Wrong' or 'very wrong' for you to take one or two drinks of an alcoholic beverage nearly every day	78.4%	86.8%	84.9%
'Wrong' or 'very wrong' for you to smoke marijuana once or twice a week	83.5%	81.1%	89.1%
'Wrong' or 'very wrong' to smoke e-cigarettes	80.0%	79.2%	unavailable
'Wrong' or 'very wrong' for you to smoke marijuana	83.5%	79.0%	87.1%
'Wrong' or 'very wrong' for you to have a drink of any type of alcohol	61.9%	68.1%	71.0%
Availability (% reflects very easy + sort of easy)			
'Very' or 'sort of easy' to get Cold/Cough Medicines or OTC medicines	50.5%	51.6%	47.3%
'Very' or 'sort of easy' to get alcohol	56.3%	50.7%	46.2%
'Very' or 'sort of easy' to get cigarettes	58.4%	46.0%	47.3%
'Very' or 'sort of easy' to get e-cigarettes	44.6%	43.1%	unavailable
'Very' or 'sort of easy' to get marijuana	37.2%	37.2%	29.0%
'Very' or 'sort of easy' to get Rx drugs that have not been prescribed to them	26.8%	27.7%	25.8%
'Very' or 'sort of easy' to get synthetic drugs	24.8%	22.0%	21.7%
'Very' or 'sort of easy' to get other illegal drugs	16.1%	14.0%	10.6%
Perception of Coolness (% reflects pretty cool + very coo	<u>ol)</u>		
Peers believe someone your age would be 'pretty cool' or 'very cool' if they smoked marijuana	18.1%	24.9%	16.0%
Peers believe someone your age would be 'pretty cool' or 'very cool' if they drank alcohol	25.2%	24.8%	19.6%
Peers believe someone your age would be'pretty cool' or 'very cool' if they smoked e-cigarettes	13.9%	17.0%	unavailable
Peers believe someone your age would be 'pretty cool' or 'very cool' if they smoked cigarettes	7.8%	9.1%	9.8%

Perception of Harm (% reflects slight risk + no risk at all) NOTE - % grouping is consistent with District Reports but other reports use moderate + great risk 'No risk at all' or 'slight risk' if they smoke e-cigarettes 38.6% 40.6% unavailable 'No risk at all' or 'slight risk' if they smoke marijuana 34.2% 26.9% once or twice a week 36.5% 'No risk at all' or 'slight risk' if they drink alcohol (no 35.1% 33.7% unavailable dosage specified) 'No risk at all' or 'slight risk' if they take one or two 38.6% 29.4% 37.2% drinks of an alcoholic beverage nearly every day 'No risk at all' or 'slight risk' if they use Cold/ Cough 21.7% 22.4% 18.1% Medicines or OTC medicines to get high 'No risk at all' or 'slight risk' if they have five or more 30.7% 21.5% 26.3% drinks of an alcoholic beverage once or twice a week 'No risk at all' or 'slight risk' if they smoke one or more 15.0% 14.9% 12.9% packs of cigarettes per day 'No risk at all' or 'slight risk' if they use RX drugs that have not been prescribed to them 15.8% 13.7% 10.6% 'No risk at all' or 'slight risk' if they use synthetic drugs 10.5% 11.6% 8.6% 'No risk at all' or 'slight risk' if they use any other illegal drugs or club drugs 8.7% 9.1% 6.5%



Questions have been abbreviated in order to fit within this report. For a copy of the survey which includes the full wording see https://dmh.mo.gov/ada/mobhew/

Data included in this report, excluding sample sizes, are weighted data. State level data uses a random sample. Please contact Susan Depue at susan.depue@mimh.edu with any questions.