



2016 Missouri Student Survey Morgan County



	2016 County Data	2016 MO Data*	2014 County Data
Approximate Sample Size	510		302
Demographics (% reflects "yes" answer)			
Grade level. 6th Grade	12.1%	13.9%	0.5%
Grade level. 7th Grade	14.3%	14.4%	27.8%
Grade level. 8th Grade	21.7%	14.3%	5.6%
Grade level. 9th Grade	15.4%	15.2%	23.7%
Grade level. 10th Grade	12.1%	14.7%	4.0%
Grade level. 11th Grade	13.6%	13.9%	35.4%
Grade level. 12th Grade	10.7%	13.6%	3.0%
Male	50.0%	47.7%	47.0%
Hispanic or Latino	1.1%	2.1%	1.0%
Race, alone or in combination. Black or African Am.	1.1%	16.0%	0.5%
Race, alone or in combination. White	96.3%	78.8%	98.5%
Race, alone or in combination. Other	0.7%	1.2%	0.0%

Note: Small sample sizes at the county level can make comparisons between years difficult. Take careful note of the demographic information listed above. Drastic changes in the sample, especially changes in the percent per grade, can influence the rest of the data in this report. This means changes between years may be a result of differing samples rather than an actual population change.

*Missouri data reflects a random sample, weighted to proportionally reflect the state student population.

Substances: Past 30 days (% reflects "1+" answer)			
Used alcohol	11.9%	14.2%	17.2%
Used electronic cigarettes	8.9%	10.6%	8.6%
Used Rx not prescribed for you by a doctor	10.0%	10.0%	4.1%
Used tobacco (cigarettes or chew)	12.6%	9.9%	20.2%
Used marijuana	4.1%	7.0%	9.7%
Used hookahs or water pipes	2.9%	3.4%	6.1%
Used 'OTC' drugs for non-medical reasons	0.7%	2.2%	3.6%
Used inhalants	1.1%	1.1%	1.0%
Used synthetic drugs	0.4%	0.3%	1.0%

	2016 County Data	2016 MO Data *	2014 County Data
Substances: Ever in your life (% reflects "1+" answer)			
Used hallucinogens such as LSD (acid), PCP (angel dust), or magic mushrooms	1.1%	1.1%	3.6%
Used cocaine or 'crack'	1.9%	0.9%	1.5%
Used 'club drugs' such as ecstasy	0.4%	0.8%	3.6%
Used methamphetamine (meth, crank, crystal, ice)	1.1%	0.3%	2.6%
Used heroin or 'smack'	0.4%	0.2%	1.0%
Behavior: Last 30 Days (% reflects "1+" answer)			
Missed whole day(s) of school because you skipped or cut	31.3%	29.0%	29.6%
Did not go to school because you felt you would be unsafe at school or on your way to or from school	5.2%	5.9%	7.6%
Carried a weapon such as a gun, knife or club on school property	5.5%	4.0%	6.5%
Rode with someone who was drinking alcohol	17.9%	14.3%	16.7%
Behavior: Last 3 Months (% reflects "1+" answer)			
Spread mean rumors or lies about other kids at school	22.9%	22.9%	27.9%
Posted something online or sent a text that might embarrass or hurt another student	19.9%	16.5%	16.3%
Made fun of other people	51.9%	54.1%	64.0%
Hit, shoved, or pushed another student and was not fooling around	18.1%	15.5%	17.6%
Behavior: Last 12 Months (% reflects "1+" answer)			
Been bullied on school property	37.6%	28.8%	39.4%
Were in a physical fight	19.6%	17.3%	22.7%
Been threatened or injured with a weapon such as a gun, knife or club on school property	7.4%	7.4%	9.1%
Considered suicide	12.3%	13.9%	13.1%
Planned suicide	11.2%	9.9%	8.6%
Attempted suicide	6.4%	6.3%	6.1%
Depression: Past Month (% reflects often + always)			
Was very sad	21.0%	22.7%	24.2%
Was grouchy or irritable, or in a bad mood	32.8%	33.6%	34.7%
Felt hopeless about the future	11.1%	13.5%	11.7%
Felt like not eating or eating more than usual	19.3%	21.7%	19.2%
Felt like sleeping a lot more or a lot less than usual	22.6%	24.8%	18.3%
Had difficulty concentrating on school work	27.7%	29.6%	32.1%

	2016 County Data	2016 MO Data *	2014 County Data
Perception of wrongness (% reflects wrong + very wrong)			
'Wrong' or 'very wrong' to use Rx drugs that have not been prescribed to you	93.3%	94.0%	90.4%
'Wrong' or 'very wrong' to use Cold/Cough Medicines or OTC meds to get high	94.1%	92.3%	86.7%
'Wrong' or 'very wrong' to have 5+ drinks, once or twice a week	89.1%	88.7%	79.3%
'Wrong' or 'very wrong' for you to smoke tobacco	88.2%	88.4%	80.3%
'Wrong' or 'very wrong' for you to take one or two drinks of an alcoholic beverage nearly every day	87.7%	86.8%	76.0%
'Wrong' or 'very wrong' for you to smoke marijuana once or twice a week	86.3%	81.1%	79.2%
'Wrong' or 'very wrong' to smoke e-cigarettes	85.1%	79.2%	unavailable
'Wrong' or 'very wrong' for you to smoke marijuana	85.8%	79.0%	77.4%
'Wrong' or 'very wrong' for you to have a drink of any type of alcohol	74.1%	68.1%	63.5%
Availability (% reflects very easy + sort of easy)			
'Very' or 'sort of easy' to get Cold/Cough Medicines or OTC medicines	44.1%	51.6%	60.7%
'Very' or 'sort of easy' to get alcohol	44.8%	50.7%	55.3%
'Very' or 'sort of easy' to get cigarettes	45.4%	46.0%	55.8%
'Very' or 'sort of easy' to get e-cigarettes	34.9%	43.1%	unavailable
'Very' or 'sort of easy' to get marijuana	29.5%	37.2%	38.5%
'Very' or 'sort of easy' to get Rx drugs that have not been prescribed to them	28.3%	27.7%	30.6%
'Very' or 'sort of easy' to get synthetic drugs	25.6%	22.0%	37.4%
'Very' or 'sort of easy' to get other illegal drugs	16.7%	14.0%	16.9%
Perception of Coolness (% reflects pretty cool + very cool)			
Peers believe someone your age would be 'pretty cool' or 'very cool' if they smoked marijuana	18.2%	24.9%	22.2%
Peers believe someone your age would be 'pretty cool' or 'very cool' if they drank alcohol	20.7%	24.8%	25.6%
Peers believe someone your age would be 'pretty cool' or 'very cool' if they smoked e-cigarettes	13.4%	17.0%	unavailable
Peers believe someone your age would be 'pretty cool' or 'very cool' if they smoked cigarettes	6.7%	9.1%	10.7%

Perception of Harm (% reflects slight risk + no risk at all)

NOTE - % grouping is consistent with District Reports but other reports use moderate + great risk

'No risk at all' or 'slight risk' if they smoke e-cigarettes	31.3%	40.6%	unavailable
'No risk at all' or 'slight risk' if they smoke marijuana once or twice a week	29.4%	36.5%	40.4%
'No risk at all' or 'slight risk' if they drink alcohol (no dosage specified)	29.0%	33.7%	unavailable
'No risk at all' or 'slight risk' if they take one or two drinks of an alcoholic beverage nearly every day	32.3%	29.4%	46.7%
'No risk at all' or 'slight risk' if they use Cold/ Cough Medicines or OTC medicines to get high	17.8%	22.4%	27.6%
'No risk at all' or 'slight risk' if they have five or more drinks of an alcoholic beverage once or twice a week	21.6%	21.5%	36.7%
'No risk at all' or 'slight risk' if they smoke one or more packs of cigarettes per day	12.1%	14.9%	17.9%
'No risk at all' or 'slight risk' if they use RX drugs that have not been prescribed to them	11.9%	13.7%	17.8%
'No risk at all' or 'slight risk' if they use synthetic drugs	9.4%	11.6%	11.9%
'No risk at all' or 'slight risk' if they use any other illegal drugs or club drugs	7.1%	9.1%	7.7%



Questions have been abbreviated in order to fit within this report. For a copy of the survey which includes the full wording see <https://dmh.mo.gov/ada/mobhew/>
 Data included in this report, excluding sample sizes, are weighted data. State level data uses a random sample. Please contact Susan Depue at susan.depue@mimh.edu with any questions.