

2016 Missouri Student Survey Shannon County





2016	2016	2014
County Data	MO Data*	County Data

Approximate Sample Size	138		47
Demographics (% reflects "yes" answer)			
Grade level. 6th Grade	18.6%	13.9%	30.0%
Grade level. 7th Grade	13.3%	14.4%	16.3%
Grade level. 8th Grade	19.5%	14.3%	0.0%
Grade level. 9th Grade	12.4%	15.2%	52.5%
Grade level. 10th Grade	15.9%	14.7%	1.3%
Grade level. 11th Grade	9.7%	13.9%	0.0%
Grade level. 12th Grade	10.6%	13.6%	0.0%
Male	49.6%	47.7%	38.3%
Hispanic or Latino	1.9%	2.1%	1.2%
Race, alone or in combination. Black or African Am.	0.0%	16.0%	0.0%
Race, alone or in combination. White	96.4%	78.8%	100.0%
Race, alone or in combination. Other	0.0%	1.2%	0.0%

Note: Small sample sizes at the county level can make comparisons between years difficult. Take careful note of the demographic information listed above. Drastic changes in the sample, especially changes in the percent per grade, can influence the rest of the data in this report. This means changes between years may be a result of differing samples rather than an actual population change.

^{*}Missouri data reflects a random sample, weighted to proportionally reflect the state student population.

Substances: Past 30 days (% reflects "1+" answer)			
Used alcohol	17.0%	14.2%	16.3%
Used electronic cigarettes	10.6%	10.6%	14.9%
Used Rx not prescribed for you by a doctor	8.0%	10.0%	12.5%
Used tobacco (cigarettes or chew)	20.5%	9.9%	40.0%
Used marijuana	8.8%	7.0%	17.7%
Used hookahs or water pipes	4.5%	3.4%	7.5%
Used 'OTC' drugs for non-medical reasons	0.9%	2.2%	0.0%
Used inhalants	0.9%	1.1%	0.0%
Used synthetic drugs	1.8%	0.3%	5.1%

	2016 County Data	2016 MO Data *	2014 County Data
Substances: Ever in your life (% reflects "1+" answer)			
Used hallucinogens such as LSD (acid), PCP (angel dust), or magic mushrooms	2.7%	1.1%	5.0%
Used cocaine or 'crack'	0.9%	0.9%	0.0%
Used 'club drugs' such as ecstasy	0.9%	0.8%	7.4%
Used methamphetamine (meth, crank, crystal, ice)	1.8%	0.3%	5.0%
Used heroin or 'smack'	0.0%	0.2%	0.0%
Behavior: Last 30 Days (% reflects "1+" answer)			
Missed whole day(s) of school because you skipped or cut	44.7%	29.0%	51.3%
Did not go to school because you felt you would be unsafe at school or on your way to or from school	5.4%	5.9%	1.3%
Carried a weapon such as a gun, knife or club on school property	11.6%	4.0%	19.8%
Rode with someone who was drinking alcohol	12.5%	14.3%	7.4%
Behavior: Last 3 Months (% reflects "1+" answer)			
Spread mean rumors or lies about other kids at school	13.3%	22.9%	26.6%
Posted something online or sent a text that might embarrass or hurt another student	13.4%	16.5%	11.4%
Made fun of other people	41.6%	54.1%	36.7%
Hit, shoved, or pushed another student and was not fooling around	22.5%	15.5%	31.3%
Behavior: Last 12 Months (% reflects "1+" answer)			
Been bullied on school property	24.3%	28.8%	30.7%
Were in a physical fight	19.5%	17.3%	24.1%
Been threatened or injured with a weapon such as a gun, knife or club on school property	8.0%	7.4%	11.8%
Considered suicide	13.6%	13.9%	10.0%
Planned suicide	10.3%	9.9%	10.7%
Attempted suicide	5.5%	6.3%	1.3%
<u>Depression: Past Month (% reflects often + always)</u>			
Was very sad	22.3%	22.7%	34.2%
Was grouchy or irritable, or in a bad mood	28.6%	33.6%	19.8%
Felt hopeless about the future	10.7%	13.5%	2.5%
Felt like not eating or eating more than usual	18.9%	21.7%	3.8%
Felt like sleeping a lot more or a lot less than usual	28.3%	24.8%	11.1%
Had difficulty concentrating on school work	28.3%	29.6%	8.6%

	2016 County Data	2016 MO Data *	2014 County Data
Perception of wrongness (% reflects wrong + very wrong))		
'Wrong' or 'very wrong' to use Rx drugs that have not been prescribed to you	93.8%	94.0%	91.3%
'Wrong' or 'very wrong' to use Cold/Cough Medicines or OTC meds to get high	93.8%	92.3%	92.6%
'Wrong' or 'very wrong' to have 5+ drinks, once or twice a week	82.0%	88.7%	78.7%
'Wrong' or 'very wrong' for you to smoke tobacco	84.8%	88.4%	53.1%
'Wrong' or 'very wrong' for you to take one or two drinks of an alcoholic beverage nearly every day	82.9%	86.8%	79.0%
'Wrong' or 'very wrong' for you to smoke marijuana once or twice a week	83.0%	81.1%	76.3%
'Wrong' or 'very wrong' to smoke e-cigarettes	78.6%	79.2%	unavailable
'Wrong' or 'very wrong' for you to smoke marijuana	81.8%	79.0%	76.5%
'Wrong' or 'very wrong' for you to have a drink of any type of alcohol	64.3%	68.1%	63.0%
Availability (% reflects very easy + sort of easy)			
'Very' or 'sort of easy' to get Cold/Cough Medicines or OTC medicines	51.4%	51.6%	34.6%
'Very' or 'sort of easy' to get alcohol	42.7%	50.7%	38.8%
'Very' or 'sort of easy' to get cigarettes	47.3%	46.0%	45.0%
'Very' or 'sort of easy' to get e-cigarettes	28.2%	43.1%	unavailable
'Very' or 'sort of easy' to get marijuana	36.9%	37.2%	29.6%
'Very' or 'sort of easy' to get Rx drugs that have not been prescribed to them	27.9%	27.7%	25.0%
'Very' or 'sort of easy' to get synthetic drugs	23.2%	22.0%	13.8%
'Very' or 'sort of easy' to get other illegal drugs	13.4%	14.0%	14.7%
Perception of Coolness (% reflects pretty cool + very coo	<u>ol)</u>		
Peers believe someone your age would be 'pretty cool' or 'very cool' if they smoked marijuana	21.4%	24.9%	38.8%
Peers believe someone your age would be 'pretty cool' or 'very cool' if they drank alcohol	26.5%	24.8%	33.8%
Peers believe someone your age would be'pretty cool' or 'very cool' if they smoked e-cigarettes	9.0%	17.0%	unavailable
Peers believe someone your age would be 'pretty cool' or 'very cool' if they smoked cigarettes	7.1%	9.1%	27.5%

10.0%

Perception of Harm (% reflects slight risk + no risk at all) NOTE - % grouping is consistent with District Reports but other reports use moderate + great risk 'No risk at all' or 'slight risk' if they smoke e-cigarettes 54.6% 40.6% unavailable 'No risk at all' or 'slight risk' if they smoke marijuana once or twice a week 38.4% 36.5% 50.6% 'No risk at all' or 'slight risk' if they drink alcohol (no 40.7% 33.7% unavailable dosage specified) 'No risk at all' or 'slight risk' if they take one or two 50.4% 29.4% 57.5% drinks of an alcoholic beverage nearly every day 'No risk at all' or 'slight risk' if they use Cold/ Cough 22.1% 22.4% 16.2% Medicines or OTC medicines to get high 'No risk at all' or 'slight risk' if they have five or more 36.9% 21.5% 47.5% drinks of an alcoholic beverage once or twice a week 'No risk at all' or 'slight risk' if they smoke one or more 27.4% 14.9% 24.7% packs of cigarettes per day 'No risk at all' or 'slight risk' if they use RX drugs that have not been prescribed to them 14.2% 13.7% 22.5% 'No risk at all' or 'slight risk' if they use synthetic drugs 14.3% 11.6% 12.5% 'No risk at all' or 'slight risk' if they use any other illegal



Questions have been abbreviated in order to fit within this report. For a copy of the survey which includes the full wording see https://dmh.mo.gov/ada/mobhew/

Data included in this report, excluding sample sizes, are weighted data. State level data uses a random sample. Please contact Susan Depue at susan.depue@mimh.edu with any questions.