



2016 Missouri Student Survey Taney County



	2016 County Data	2016 MO Data*	2014 County Data
Approximate Sample Size	601		1246
Demographics (% reflects "yes" answer)			
Grade level. 6th Grade	7.1%	13.9%	19.2%
Grade level. 7th Grade	14.2%	14.4%	4.3%
Grade level. 8th Grade	24.4%	14.3%	22.9%
Grade level. 9th Grade	22.1%	15.2%	31.6%
Grade level. 10th Grade	18.9%	14.7%	2.2%
Grade level. 11th Grade	8.2%	13.9%	7.6%
Grade level. 12th Grade	5.0%	13.6%	12.1%
Male	48.1%	47.7%	47.8%
Hispanic or Latino	3.5%	2.1%	2.7%
Race, alone or in combination. Black or African Am.	0.7%	16.0%	1.0%
Race, alone or in combination. White	93.1%	78.8%	97.9%
Race, alone or in combination. Other	1.0%	1.2%	1.3%

Note: Small sample sizes at the county level can make comparisons between years difficult. Take careful note of the demographic information listed above. Drastic changes in the sample, especially changes in the percent per grade, can influence the rest of the data in this report. This means changes between years may be a result of differing samples rather than an actual population change.

*Missouri data reflects a random sample, weighted to proportionally reflect the state student population.

Substances: Past 30 days (% reflects "1+" answer)			
Used alcohol	13.3%	14.2%	13.2%
Used electronic cigarettes	15.2%	10.6%	13.8%
Used Rx not prescribed for you by a doctor	8.0%	10.0%	3.2%
Used tobacco (cigarettes or chew)	11.1%	9.9%	12.7%
Used marijuana	7.9%	7.0%	7.8%
Used hookahs or water pipes	4.6%	3.4%	8.3%
Used 'OTC' drugs for non-medical reasons	1.0%	2.2%	1.8%
Used inhalants	2.4%	1.1%	1.0%
Used synthetic drugs	0.4%	0.3%	0.7%

	2016 County Data	2016 MO Data *	2014 County Data
Substances: Ever in your life (% reflects "1+" answer)			
Used hallucinogens such as LSD (acid), PCP (angel dust), or magic mushrooms	2.2%	1.1%	2.4%
Used cocaine or 'crack'	1.6%	0.9%	1.3%
Used 'club drugs' such as ecstasy	1.7%	0.8%	2.3%
Used methamphetamine (meth, crank, crystal, ice)	0.8%	0.3%	1.0%
Used heroin or 'smack'	0.8%	0.2%	0.7%
Behavior: Last 30 Days (% reflects "1+" answer)			
Missed whole day(s) of school because you skipped or cut	31.2%	29.0%	28.0%
Did not go to school because you felt you would be unsafe at school or on your way to or from school	7.0%	5.9%	5.4%
Carried a weapon such as a gun, knife or club on school property	6.9%	4.0%	5.9%
Rode with someone who was drinking alcohol	14.0%	14.3%	11.4%
Behavior: Last 3 Months (% reflects "1+" answer)			
Spread mean rumors or lies about other kids at school	27.9%	22.9%	23.2%
Posted something online or sent a text that might embarrass or hurt another student	14.5%	16.5%	17.4%
Made fun of other people	58.7%	54.1%	52.6%
Hit, shoved, or pushed another student and was not fooling around	16.8%	15.5%	17.8%
Behavior: Last 12 Months (% reflects "1+" answer)			
Been bullied on school property	43.1%	28.8%	39.6%
Were in a physical fight	19.3%	17.3%	17.8%
Been threatened or injured with a weapon such as a gun, knife or club on school property	9.8%	7.4%	7.6%
Considered suicide	20.7%	13.9%	14.3%
Planned suicide	13.9%	9.9%	10.5%
Attempted suicide	7.7%	6.3%	5.9%
Depression: Past Month (% reflects often + always)			
Was very sad	31.3%	22.7%	23.1%
Was grouchy or irritable, or in a bad mood	34.4%	33.6%	25.6%
Felt hopeless about the future	17.5%	13.5%	12.4%
Felt like not eating or eating more than usual	23.0%	21.7%	19.3%
Felt like sleeping a lot more or a lot less than usual	28.1%	24.8%	20.8%
Had difficulty concentrating on school work	33.9%	29.6%	25.9%

	2016 County Data	2016 MO Data *	2014 County Data
Perception of wrongness (% reflects wrong + very wrong)			
'Wrong' or 'very wrong' to use Rx drugs that have not been prescribed to you	94.4%	94.0%	94.2%
'Wrong' or 'very wrong' to use Cold/Cough Medicines or OTC meds to get high	93.7%	92.3%	93.9%
'Wrong' or 'very wrong' to have 5+ drinks, once or twice a week	88.5%	88.7%	87.8%
'Wrong' or 'very wrong' for you to smoke tobacco	86.3%	88.4%	85.3%
'Wrong' or 'very wrong' for you to take one or two drinks of an alcoholic beverage nearly every day	86.5%	86.8%	85.3%
'Wrong' or 'very wrong' for you to smoke marijuana once or twice a week	81.7%	81.1%	81.5%
'Wrong' or 'very wrong' to smoke e-cigarettes	77.4%	79.2%	unavailable
'Wrong' or 'very wrong' for you to smoke marijuana	80.8%	79.0%	79.0%
'Wrong' or 'very wrong' for you to have a drink of any type of alcohol	72.7%	68.1%	69.4%
Availability (% reflects very easy + sort of easy)			
'Very' or 'sort of easy' to get Cold/Cough Medicines or OTC medicines	53.2%	51.6%	53.0%
'Very' or 'sort of easy' to get alcohol	51.3%	50.7%	48.2%
'Very' or 'sort of easy' to get cigarettes	47.5%	46.0%	49.0%
'Very' or 'sort of easy' to get e-cigarettes	47.1%	43.1%	unavailable
'Very' or 'sort of easy' to get marijuana	39.8%	37.2%	35.6%
'Very' or 'sort of easy' to get Rx drugs that have not been prescribed to them	30.1%	27.7%	27.4%
'Very' or 'sort of easy' to get synthetic drugs	27.0%	22.0%	24.2%
'Very' or 'sort of easy' to get other illegal drugs	14.5%	14.0%	16.6%
Perception of Coolness (% reflects pretty cool + very cool)			
Peers believe someone your age would be 'pretty cool' or 'very cool' if they smoked marijuana	24.3%	24.9%	23.1%
Peers believe someone your age would be 'pretty cool' or 'very cool' if they drank alcohol	19.6%	24.8%	21.3%
Peers believe someone your age would be 'pretty cool' or 'very cool' if they smoked e-cigarettes	21.7%	17.0%	unavailable
Peers believe someone your age would be 'pretty cool' or 'very cool' if they smoked cigarettes	9.2%	9.1%	11.0%

Perception of Harm (% reflects slight risk + no risk at all)

NOTE - % grouping is consistent with District Reports but other reports use moderate + great risk

'No risk at all' or 'slight risk' if they smoke e-cigarettes	45.6%	40.6%	unavailable
'No risk at all' or 'slight risk' if they smoke marijuana once or twice a week	35.8%	36.5%	37.3%
'No risk at all' or 'slight risk' if they drink alcohol (no dosage specified)	30.8%	33.7%	unavailable
'No risk at all' or 'slight risk' if they take one or two drinks of an alcoholic beverage nearly every day	35.6%	29.4%	40.3%
'No risk at all' or 'slight risk' if they use Cold/ Cough Medicines or OTC medicines to get high	20.8%	22.4%	17.7%
'No risk at all' or 'slight risk' if they have five or more drinks of an alcoholic beverage once or twice a week	23.7%	21.5%	27.0%
'No risk at all' or 'slight risk' if they smoke one or more packs of cigarettes per day	14.4%	14.9%	11.8%
'No risk at all' or 'slight risk' if they use RX drugs that have not been prescribed to them	11.1%	13.7%	12.8%
'No risk at all' or 'slight risk' if they use synthetic drugs	10.0%	11.6%	7.5%
'No risk at all' or 'slight risk' if they use any other illegal drugs or club drugs	6.8%	9.1%	5.9%



Questions have been abbreviated in order to fit within this report. For a copy of the survey which includes the full wording see <https://dmh.mo.gov/ada/mobhew/>
 Data included in this report, excluding sample sizes, are weighted data. State level data uses a random sample. Please contact Susan Depue at susan.depue@mimh.edu with any questions.