

Approximate Sample Size

Grade level. 6th Grade

Grade level. 7th Grade

Grade level. 8th Grade

Grade level. 9th Grade Grade level. 10th Grade

Grade level. 11th Grade

Grade level. 12th Grade

Hispanic or Latino

Male

Demographics (% reflects "yes" answer)

Race, alone or in combination. Black or African Am.

Race, alone or in combination. White

Race, alone or in combination. Other

## 2016 Missouri Student Survey New Madrid County





2016 County Data	2016 MO Data*	2014 County Data
319		369
12.4%	13.9%	11.0%
13.4%	14.4%	17.7%
18.3%	14.3%	17.2%
16.6%	15.2%	14.8%
14.1%	14.7%	19.1%
14.5%	13.9%	12.0%
10.7%	13.6%	8.1%
47.6%	47.7%	47.4%
1.0%	2.1%	1.0%
14.3%	16.0%	16.0%

78.8%

1.2%

84.0%

0.5%

83.6%

0.7%

Note: Small sample sizes at the county level can make comparisons between years difficult. Take careful note of the demographic information listed above. Drastic changes in the sample, especially changes in the percent per grade, can influence the rest of the data in this report. This means changes between years may be a result of differing samples rather than an actual population change.

<sup>\*</sup>Missouri data reflects a random sample, weighted to proportionally reflect the state student population.

Substances: Past 30 days (% reflects "1+" answer)					
Used alcohol	16.7%	14.2%	18.8%		
Used electronic cigarettes	12.6%	10.6%	8.6%		
Used Rx not prescribed for you by a doctor	11.9%	10.0%	2.9%		
Used tobacco (cigarettes or chew)	12.3%	9.9%	18.8%		
Used marijuana	4.1%	7.0%	6.8%		
Used hookahs or water pipes	2.4%	3.4%	2.4%		
Used 'OTC' drugs for non-medical reasons	1.7%	2.2%	2.4%		
Used inhalants	1.7%	1.1%	2.4%		
Used synthetic drugs	0.0%	0.3%	1.0%		

	2016 County Data	2016 MO Data *	2014 County Data
Substances: Ever in your life (% reflects "1+" answer)			
Used hallucinogens such as LSD (acid), PCP (angel dust), or magic mushrooms	0.3%	1.1%	2.0%
Used cocaine or 'crack'	0.0%	0.9%	1.4%
Used 'club drugs' such as ecstasy	1.0%	0.8%	0.0%
Used methamphetamine (meth, crank, crystal, ice)	1.0%	0.3%	1.4%
Used heroin or 'smack'	0.0%	0.2%	0.5%
Behavior: Last 30 Days (% reflects "1+" answer)			
Missed whole day(s) of school because you skipped or cut	31.0%	29.0%	34.6%
Did not go to school because you felt you would be unsafe at school or on your way to or from school	2.4%	5.9%	3.3%
Carried a weapon such as a gun, knife or club on school property	2.4%	4.0%	4.8%
Rode with someone who was drinking alcohol	21.2%	14.3%	21.1%
Behavior: Last 3 Months (% reflects "1+" answer)			
Spread mean rumors or lies about other kids at school	25.5%	22.9%	29.2%
Posted something online or sent a text that might embarrass or hurt another student	16.0%	16.5%	21.2%
Made fun of other people	56.2%	54.1%	63.0%
Hit, shoved, or pushed another student and was not fooling around	20.3%	15.5%	23.2%
Behavior: Last 12 Months (% reflects "1+" answer)			
Been bullied on school property	27.6%	28.8%	26.0%
Were in a physical fight	20.4%	17.3%	23.8%
Been threatened or injured with a weapon such as a gun, knife or club on school property	5.8%	7.4%	6.7%
Considered suicide	9.5%	13.9%	15.2%
Planned suicide	5.5%	9.9%	8.6%
Attempted suicide	3.7%	6.3%	6.7%
Depression: Past Month (% reflects often + always)			
Was very sad	17.0%	22.7%	18.8%
Was grouchy or irritable, or in a bad mood	32.2%	33.6%	25.6%
Felt hopeless about the future	7.2%	13.5%	11.5%
Felt like not eating or eating more than usual	16.7%	21.7%	18.4%
Felt like sleeping a lot more or a lot less than usual	18.8%	24.8%	16.1%
Had difficulty concentrating on school work	24.8%	29.6%	18.8%

	2016 County Data	2016 MO Data *	2014 County Data
Perception of wrongness (% reflects wrong + very wrong	)		
'Wrong' or 'very wrong' to use Rx drugs that have not been prescribed to you	95.2%	94.0%	93.2%
'Wrong' or 'very wrong' to use Cold/Cough Medicines or OTC meds to get high	94.2%	92.3%	92.8%
'Wrong' or 'very wrong' to have 5+ drinks, once or twice a week	84.9%	88.7%	76.1%
'Wrong' or 'very wrong' for you to smoke tobacco	85.7%	88.4%	76.9%
'Wrong' or 'very wrong' for you to take one or two drinks of an alcoholic beverage nearly every day	82.5%	86.8%	70.5%
'Wrong' or 'very wrong' for you to smoke marijuana once or twice a week	89.1%	81.1%	84.6%
'Wrong' or 'very wrong' to smoke e-cigarettes	77.8%	79.2%	unavailable
'Wrong' or 'very wrong' for you to smoke marijuana	88.4%	79.0%	82.2%
'Wrong' or 'very wrong' for you to have a drink of any type of alcohol	68.9%	68.1%	58.7%
Availability (% reflects very easy + sort of easy)			
'Very' or 'sort of easy' to get Cold/Cough Medicines or OTC medicines	44.9%	51.6%	46.9%
'Very' or 'sort of easy' to get alcohol	45.7%	50.7%	53.4%
'Very' or 'sort of easy' to get cigarettes	42.5%	46.0%	55.8%
'Very' or 'sort of easy' to get e-cigarettes	38.2%	43.1%	unavailable
'Very' or 'sort of easy' to get marijuana	18.4%	37.2%	31.6%
'Very' or 'sort of easy' to get Rx drugs that have not been prescribed to them	22.6%	27.7%	27.4%
'Very' or 'sort of easy' to get synthetic drugs	12.9%	22.0%	16.3%
'Very' or 'sort of easy' to get other illegal drugs	7.5%	14.0%	12.1%
Perception of Coolness (% reflects pretty cool + very cool	<u>ol)</u>		
Peers believe someone your age would be 'pretty cool' or 'very cool' if they smoked marijuana	12.6%	24.9%	20.6%
Peers believe someone your age would be 'pretty cool' or 'very cool' if they drank alcohol	23.2%	24.8%	29.7%
Peers believe someone your age would be'pretty cool' or 'very cool' if they smoked e-cigarettes	14.6%	17.0%	unavailable
Peers believe someone your age would be 'pretty cool' or 'very cool' if they smoked cigarettes	7.5%	9.1%	15.9%

## Perception of Harm (% reflects slight risk + no risk at all) NOTE - % grouping is consistent with District Reports but other reports use moderate + great risk 'No risk at all' or 'slight risk' if they smoke e-cigarettes 46.7% 40.6% unavailable 'No risk at all' or 'slight risk' if they smoke marijuana 32.7% once or twice a week 25.1% 36.5% 'No risk at all' or 'slight risk' if they drink alcohol (no 38.4% 33.7% unavailable dosage specified) 'No risk at all' or 'slight risk' if they take one or two 40.1% 29.4% 51.9% drinks of an alcoholic beverage nearly every day 'No risk at all' or 'slight risk' if they use Cold/ Cough 22.1% 22.4% 17.6% Medicines or OTC medicines to get high 'No risk at all' or 'slight risk' if they have five or more 30.3% 21.5% 33.3% drinks of an alcoholic beverage once or twice a week 'No risk at all' or 'slight risk' if they smoke one or more 17.9% 14.9% 18.4% packs of cigarettes per day 'No risk at all' or 'slight risk' if they use RX drugs that have not been prescribed to them 18.6% 13.7% 13.0% 'No risk at all' or 'slight risk' if they use synthetic drugs 9.3% 11.6% 7.2% 'No risk at all' or 'slight risk' if they use any other illegal drugs or club drugs 10.6% 9.1% 7.4%



Questions have been abbreviated in order to fit within this report. For a copy of the survey which includes the full wording see https://dmh.mo.gov/ada/mobhew/

Data included in this report, excluding sample sizes, are weighted data. State level data uses a random sample. Please contact Susan Depue at susan.depue@mimh.edu with any questions.